**Amino Acid-Modified Infant Formula With Iron** 

# **Product Information:** Phenex<sup>™</sup>-1

## **Amino Acid-Modified Infant Formula With Iron**

- 1 Nutrition support of infants and toddlers with phenylketonuria (PKU).
- 1 Phenylalanine-free.
- 1 Use under medical supervision.
- 1 Phenylalanine-free to allow greater intake of intact protein.
- 1 L-tyrosine, an essential amino acid that is often deficient in infants with PKU.
- 1 L-carnitine (20 mg/100 g) and taurine (40 mg/100 g) to help supply amounts normally found in human milk and foods of animal origin.
- 1 Approximately 40% of energy as fat to help achieve acceptable formula osmolality.
- 1 6.6% of energy as linoleic acid.
- 1 Nutrient profile specifically designed for infants and toddlers.
- 1 When fed according to the Abbott Nutrition Support Protocols, provides adequate amounts of all nutrients.
- 1 Lactose-free.
- 1 Gluten-free.
- 1 Halal.

### **Safety Precautions**

- 1 Give only to infants and toddlers with proven phenylketonuria who are under medical supervision. Must be supplemented with protein and fluid in prescribed amounts to completely meet phenylalanine and water requirements.
- Not intended as a sole source of nutrition.
- Not for IV use.
- 1 Do not boil mixture or use terminal sterilization.
- 1 Never use a microwave oven to warm formula. Serious burns can result.
- Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.
- To meet the nutrient needs of the infant or child, infant formula, breast milk or additional food choices must be given to supply protein and phenylalanine requirements. These may be added to the Phenex-1 mixture or prepared separately as instructed. Depending on the specific needs of each patient, preparation will vary as prescribed by physician.



#### **Powder Unflavored:**

Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil. Less than 2% of the Following: L-Leucine, L-Tyrosine, L-Proline, L-Lysine Acetate, Calcium Phosphate, DATEM, L-Glutamine, L-Valine, L-Isoleucine, L-Arginine, Potassium Phosphate, L-Alanine, Glycine, L-Asparagine, L-Serine, L-Threonine, Sodium Citrate, Potassium Citrate, Magnesium Chloride, L-Histidine, L-Methionine, Calcium Carbonate, L-Glutamic Acid, Ascorbic Acid, L-Cystine Dihydrochloride, L-Tryptophan, L-Aspartic Acid, Choline Chloride, Taurine, m-Inositol, Ferrous Sulfate, Zinc Sulfate, Ascorbyl Palmitate, L-Carnitine, dl-Alpha-Tocopheryl Acetate, Niacinamide, Mixed Tocopherols, Calcium Pantothenate, Salt, Cupric Sulfate, Thiamine Chloride Hydrochloride, Manganese Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Beta-Carotene, Potassium Iodide, Biotin, Phylloquinone, Sodium Selenate, Chromium Chloride, Sodium Molybdate, Vitamin D3, and Cyanocobalamin.

#### **Availability**

| List Number | Item   |
|-------------|--|
| 51120       | Phenex-1 Powder / 14.1-oz (400-g) Can / 6 ct |





#### **Nutrition Information - Powder Unflavored**

|                                   | 100 Cal                                    | 100 g Powder                               |  |
|-----------------------------------|--|--|--|
|                                   | Value                                      | Value                                      |  |
|                                   |  |  |  |
| Protein Equivalent Source         | L-Amino Acids                              | L-Amino Acids                              |  |
| Fat Source Source                 | High Oleic Safflower, Coconut and Soy Oils | High Oleic Safflower, Coconut and Soy Oils |  |
| Carbohydrate Source               | Corn Syrup Solids                          | Corn Syrup Solids                          |  |
| Osmolality, mOsm/kg H2O           | 370  | 370  |  |
| Protein Equivalent, g             | 3.1  | 15.0                                       |  |
| Fat, g                            | 4.5  | 21.7                                       |  |
| Carbohydrate, g                   | 11.0                                       | 53.0                                       |  |
| Linoleic Acid, mg                 | 729  | 3500                                       |  |
| Linolenic Acid, mg                | 73   | 350  |  |
| L-Carnitine, mg                   | 4  | 20   |  |
| Calories                          | 100  | 480  |  |
| Potential Renal Solute Load, mOsm | 27.8                                       | 133  |  |
| Vitamin A, IU                     | 292  | 1400                                       |  |
| Vitamin A, mcg RE                 | 87.5                                       | 420  |  |
| Vitamin D, IU                     | 63   | 300  |  |
| Vitamin D, mcg                    | 1.6  | 7.5  |  |
| Vitamin E, IU                     | 3  | 15   |  |
| Vitamin E, mg alpha-TE            | 2.1  | 10.1                                       |  |
| Vitamin K, mcg                    | 10.4                                       | 50   |  |
| Thiamin (Vitamin B1), mcg         | 396  | 1900                                       |  |
| Riboflavin (Vitamin B2), mcg      | 188  | 900  |  |
| Vitamin B6, mcg                   | 156  | 750  |  |
| Vitamin B12, mcg                  | 1.0  | 4.9  |  |
| Niacin, mcg                       | 2083                                       | 10000                                      |  |
| Niacin, mg NE                     | 2.7  | 12.8                                       |  |
| Folic Acid (Folacin), mcg         | 48   | 230  |  |
| Pantothenic Acid, mcg             | 1438                                       | 6900                                       |  |
| Biotin, mcg                       | 13.5                                       | 65   |  |
| Vitamin C, mg                     | 10.4                                       | 50   |  |
| Choline, mg                       | 16.7                                       | 80   |  |
| Inositol, mg                      | 8.3  | 40   |  |
| Calcium, mg                       | 120  | 575  |  |
| Calcium, mEq                      | 6.0  | 28.8                                       |  |
| Phosphorus, mg                    | 83   | 400  |  |
| Magnesium, mg                     | 10   | 50   |  |
| Iron, mg                          | 1.9  | 9  |  |
| Zinc, mg                          | 1.7  | 8  |  |
| Manganese, mcg                    | 104  | 500  |  |
| Copper, mcg                       | 229  | 1100                                       |  |



### **Amino Acid-Modified Infant Formula With Iron**

#### **Nutrition Information - Powder Unflavored**

|                 | 100 Cal | 100 g Powder |
|-----------------|---------|--------------|
|                 | Value   | Value        |
| lodine, mcg     | 13.5    | 65           |
| Selenium, mcg   | 4.2     | 20           |
| Chromium, mcg   | 2.3     | 11           |
| Molybdenum, mcg | 2.5     | 12           |
| Sodium, mg      | 39.6    | 190          |
| Sodium, mEq     | 1.7     | 8.3          |
| Potassium, mg   | 140.6   | 675          |
| Potassium, mEq  | 3.6     | 17.3         |
| Chloride, mg    | 67.7    | 325          |
| Chloride, mEq   | 1.9     | 9.2          |

#### **Powder Unflavored Footnotes & References**

Per 100 Cal

\*Prepared at 20 Cal/fl oz.

Per 100 g Powder

\*Prepared at 20 Cal/fl oz.

#### **Preparation**

Approximate Weights for Unpacked, Level U.S. Standard Dry Household Measures for Powder\*

| Household Measure (US) | Weight (g) |
|------------------------|------------|
| 1 Tbsp                 | 8          |
| 1/4 Cup                | 30         |
| 1/3 Cup                | 40         |
| 1/2 Cup                | 60         |
| 1 Cup                  | 120        |

<sup>\*</sup> For most accurate results, Phenex-1 should be weighed on a scale that reads in grams.

#### Directions for Preparation and Use: Use Only As Directed by a Physician

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- 1 Wash your hands, surfaces and utensils.
- 1 Pour prescribed amount of water into a clean container.
- 1 Add prescribed amount of Phenex-1 powder (and other ingredients if recommended).
- 1 Shake well for 10-15 seconds; if using a blender, mix no more than 5 seconds.
- 1 Pour mixture into clean feeding bottles or container; cap.
- 1 Once feeding begins, use within 1 hour or discard.



### **Amino Acid-Modified Infant Formula With Iron**

## **Storage & Handling**

#### Powder

- 1 Once mixed, store bottles or container in refrigerator and feed to baby or child within 24 hours
- 1 Store unopened or opened can at room temperature; avoid extreme temperatures.
- 1 Use opened can contents within 1 month.

