

Product Information: PediaSure® Peptide 1.5 Cal

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

PediaSure® Peptide 1.5 Cal

Peptide-Based Nutrition for Children

- 1 PediaSure Peptide 1.5 is a nutritionally complete, peptide-based formula for the nutritional needs of children 1-13 years with malabsorption, maldigestion, and other GI conditions. It is designed to meet the nutritional and caloric needs of kids who require higher caloric density, have fluid restrictions, and are at risk for malnutrition.
- 1 For oral or tube feeding.
- 1 For supplemental or sole-source nutrition.
- 1 Use under medical supervision.
- 1 Nutrient blend designed to help support tolerance and absorption:
 - Hydrolyzed whey-dominant protein.
 - Nutraflora® scFOS®* prebiotic.
 - Structured lipids.
- 1 Nutrition to help support:
 - Absorption.
 - Tolerance.
 - Growth and development.
- 1 Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals for children 1 to 8 years of age in 667 mL (1000 Cal); for children 9 to 13 years of age in 1000 mL (1500 Cal).
- 1 Only peptide-based pediatric product with Kosher status.
- 1 Halal.
- 1 Kosher.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.



Safety Precautions

- 1 Not intended for infants under 1 year of age unless specified by a physician.
- 1 Not for IV use.
- 1 Not for children with galactosemia.
- 1 **Feeding sets are for single-patient use only. Use clean technique to avoid set and/or product contamination.**

* NutraFlora® and scFOS® are not registered trademarks of Abbott Laboratories.

Ingredients

Liquid Vanilla:

Water, Corn Maltodextrin, Structured Lipid (Interesterified Canola, Medium-Chain Triglycerides), Whey Protein Hydrolysate, Hydrolyzed Sodium Caseinate, Medium-Chain Triglycerides, Canola Oil, Short-Chain Fructooligosaccharides. Less than 0.5% of the Following: Calcium Phosphate, Artificial Flavor, Potassium Citrate, Magnesium Phosphate, Potassium Chloride, Soy Lecithin, Ascorbic Acid, Cellulose Gel, Choline Chloride, Sodium Citrate, m-Inositol, Cellulose Gum, Taurine, Magnesium Chloride, Acesulfame K, Ferrous Sulfate, Zinc Sulfate, Sucralose, dl-Alpha-Tocopheryl Acetate, L-Carnitine, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Cupric Sulfate, Vitamin A Palmitate, Folic Acid, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate, Sodium Molybdate, Phylloquinone, Cyanocobalamin, and Vitamin D3.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
56655	PediaSure Peptide 1.5 Ready-to-Drink Vanilla / 8-fl-oz (237-mL) Bottle / 24 ct
62731	PediaSure Peptide 1.5 Ready-To-Hang / 1000-mL (1.1-QT) Bottle / 8 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)		1000 mL	
	Value	%RDI / %DV	Value	%RDI / %DV
Nutrient Density, Cal/mL	1.5		1.5	
Protein, % Cal	12		12	
Carbohydrate, % Cal	53		53	
Fat, % Cal	35		35	
MCT:LCT	50:50		50:50	
n6:n3	2.6:1		2.6:1	
mL to Meet Or Exceed 100% DRIs for Children 1-8 Years	667		667	
mL to Meet Or Exceed 100% DRIs for Children 9-13 Years	1000		1000	
Total Cal:g Nitrogen	208:1		208:1	
Nonprotein Cal:g Nitrogen	183:1		183:1	
Osmolality, mOsm/kg H2O	450		450	
Renal Solute Load, mOsm/L	439		439	
Viscosity	Thin (room temperature), Nectar (chilled)		Thin (room temperature), Nectar (chilled)	
Minimum Tube Size for Gravity/Pump Feeding (Fr)	10/8		10/8	
Protein, g	10.7		45.1	
Fat, g	14.4		60.8	
Trans Fat, g	0.0		0.0	
Carbohydrate, g	47.6		201	
Dietary Fiber, g	1.1		4.6	
scFOS, g	1.1		4.6	
L-Carnitine, mg	6.0		25	
Taurine, mg	26		110	
Water, g	182		768	
Calories	356		1500	
Vitamin A, IU	889		3750	
Vitamin D, IU	188		791	
Vitamin E, IU	8.1		34	
Vitamin K, mcg	21		89	
Vitamin C, mg	36		152	
Folic Acid, mcg	107		450	
Thiamin (Vitamin B1), mg	1.0		4.2	
Riboflavin (Vitamin B2), mg	0.8		3.3	
Vitamin B6, mg	1.0		4.2	
Vitamin B12, mcg	2.1		8.9	
Niacin, mg	3.6		15	
Choline, mg	107		450	
Biotin, mcg	68		286	
Pantothenic Acid, mg	3.6		15	

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)		1000 mL	
	Value	%RDI / %DV	Value	%RDI / %DV
m-Inositol, mg	29		122	
Sodium, mg	255		1075	
Sodium, mEq	11.1		46.7	
Potassium, mg	480		2025	
Potassium, mEq	12.3		51.9	
Chloride, mg	360		1520	
Chloride, mEq	10.1		42.8	
Calcium, mg	375		1580	
Phosphorus, mg	300		1265	
Magnesium, mg	71		300	
Iodine, mcg	35		148	
Manganese, mg	0.5		2.2	
Copper, mg	0.4		1.6	
Zinc, mg	4.3		18	
Iron, mg	5.0		21	
Selenium, mcg	11		46	
Chromium, mcg	11		46	
Molybdenum, mcg	13		54	

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Preparation

Instructions for Use: 8-fl-oz Bottle

- 1 **Shake very well.**
- 1 After opening, refrigerate and **use within 48 hours.**
- 1 Store unopened at room temperature.

Tube Feeding:

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to child's condition and tolerance.
- 1 Additional fluid may be required.

Oral Feeding:

- 1 Best when chilled.

Instructions for Use: Ready-To-Hang Container

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination. NOTE: Failure to follow the

Instructions for Use increases the potential for microbial contamination and reduces hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Always follow directions for use provided with feeding set.

For Use with Enteral Feeding Set:

- 1 Remove the dust cover from the Abbott RTH Safety Screw Cap.
- 1 Remove the dust cover from the safety connector on the feeding set.
- 1 Insert the safety connector into the port on the Abbott RTH Safety Screw Cap, completely pierce foil.
- 1 Turn the safety connector clockwise until it is securely fastened.
- 1 Close clamp on set before inverting container.
- 1 Invert container and suspend, using the hanging ring on the bottom of the container.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.