

Product Information: Suplena® with Carb Steady®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Suplena® with Carb Steady®

For People With Chronic Kidney Disease

- 1 SUPLENA with Carb Steady is therapeutic nutrition specifically designed for chronic kidney disease (stages 3 & 4).
- 1 For tube or oral feeding.
- 1 For supplemental or sole-source nutrition.
- 1 Use under medical supervision.
- 1 Therapeutic nutrition that can help people with chronic kidney disease (stages 3 and 4) maintain their nutritional status while adhering to their renal diets.
- 1 Has Carb Steady carbohydrate blend designed to help manage blood glucose response.
- 1 High in calories —1.8 Cal/mL.
- 1 Low in protein (10% of total calories).
- 1 Low in phosphorus, potassium, calcium and sodium.
- 1 A good source of fiber, including prebiotics.
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.
- 1 Low-residue.
- 1 Suitable for lactose intolerance.



Safety Precautions

- 1 Not for patients with galactosemia.
- 1 Not for IV use.

Ingredients

Liquid Homemade Vanilla:

Water, Corn Maltodextrin, High Oleic Safflower Oil, Milk Protein Isolate, Isomaltulose, Sugar (Sucrose), Canola Oil, Sodium Caseinate, Glycerine, Short-Chain Fructooligosaccharides. Less than 1.0% of the Following: Soy Lecithin, Natural & Artificial Flavor, Potassium Citrate, Sodium Citrate, Magnesium Chloride, Choline Chloride, Magnesium Phosphate, Calcium Carbonate, Ascorbic Acid, L-Carnitine, Calcium Phosphate, Carrageenan, Potassium Chloride, Taurine, Salt (Sodium Chloride), Zinc Sulfate, d-Alpha-Tocopheryl Acetate, Ferrous Sulfate, Niacinamide, Calcium Pantothenate, Pyridoxine Hydrochloride, Manganese Sulfate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Riboflavin, Vitamin A Palmitate, Folic Acid, Chromium Chloride, Biotin, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Cyanocobalamin, and Vitamin D3.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
62088	Suplena w Carb Steady Shake Drink Homemade Vanilla Inst / 8 fl-oz (237-mL) can / 24 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Suplena® with Carb Steady®

For People With Chronic Kidney Disease

Nutrition Information - Liquid Homemade Vanilla

	8 fl oz (237 mL)	
	Value	%RDI* / %DV
Osmolality, mOsm/kg H ₂ O	780	
Renal Solute Load, mOsm/L	344	
Protein, g	10.6	
Fat, g	22.7	
Saturated Fat, g	2.0	
Trans Fat, g	0.0	
Polyunsaturated Fat, g	4.1	
Monounsaturated Fat, g	16.1	
Cholesterol, mg	5.8	
Carbohydrate, g	46.4	
Dietary Fiber, g	3.0	
Sugars, g	14.8	
Glycerine, g	2.6	
L-Carnitine, mg	63	
Taurine, mg	38	
Water, g	175	
Calories	425	
Vitamin A, IU	750	15
Vitamin D, IU	20	5
Vitamin E, IU	23	77
Vitamin K, mcg	20	25
Vitamin C, mg	25	42
Folic Acid, mcg	250	63
Vitamin B1, mg	0.56	37
Vitamin B2, mg	0.64	38
Vitamin B6, mg	2.0	100
Vitamin B12, mcg	2.3	38
Niacin, mg	7.5	38
Choline, mg	150	Not established
Biotin, mcg	120	38
Pantothenic Acid, mg	3.8	38
Sodium, mg	190	Not established
Sodium, mEq	8.3	
Potassium, mg	270	Not established
Potassium, mEq	6.9	
Chloride, mg	220	6
Chloride, mEq	6.2	
Calcium, mg	250	25
Phosphorus, mg	170	17

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Liquid Homemade Vanilla

	8 fl oz (237 mL)	
	Value	%RDI* / %DV
Magnesium, mg	50	13
Iodine, mcg	38	25
Manganese, mg	0.5	25
Copper, mg	0.5	25
Zinc, mg	6.4	43
Iron, mg	4.5	25
Selenium, mcg	18	26
Chromium, mcg	30	25
Molybdenum, mcg	19	25

Liquid Homemade Vanilla Footnotes & References

Per 8 fl oz (237 mL)

*RDI percentages for adults and children 4 or more years of age.

Storage & Handling

Instructions

- 1 Use by date on end of can.
- 1 Store unopened at room temperature.
- 1 Shake well.
- 1 Clean top of container before opening.
- 1 Once opened, cover, refrigerate and use within 48 hours.

Oral Feeding

- 1 May be used for total or supplemental nutrition.
- 1 May be fed at room temperature or chilled.

Tube Feeding

- 1 Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Pump feed at room temperature with an 8 Fr or larger tube; gravity feeding not recommended.
- 1 When used as a sole source of nutrition, additional water may be required and can be met by giving water between or after feedings or when flushing the tube.
- 1 Avoid contamination during preparation and use.