

Product Information: Jevity® 1.2 Cal

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Jevity® 1.2 Cal

High-Protein Nutrition With Fiber and NutraFlora® scFOS®



- 1 JEVITY 1.2 CAL is a high-protein, fiber-fortified formula that provides Complete, Balanced Nutrition® for long- or short-term tube feeding.
- 1 For tube feeding.
- 1 For supplemental or sole-source nutrition.
- 1 May be used for oral feeding of patients with altered taste perception.
- 1 Use under medical supervision.
- 1 Concentrated calories (1.2 Cal/mL) and high in protein (18.5% of Cal) to help tube-fed patients gain and maintain healthy weight. Protein is needed for lean body tissue maintenance and repair.
- 1 Meets or exceeds 100% of the RDI for 24 essential vitamins and minerals in 1200 Cal (1 Liter).
- 1 18 g of dietary fiber in 1 liter (10 g of a unique soluble and insoluble fiber blend and 8 g of NutraFlora® scFOS®*) supplies the minimum recommended daily fiber intake. Fiber helps maintain bowel function.
- 1 Includes 1.9 g of scFOS/8 fl oz (8 g/L and 12 g/1500 mL). scFOS are prebiotics that stimulate the growth of beneficial bacteria in the colon.
- 1 Halal.
- 1 Kosher.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.

Safety Precautions

- 1 Not for IV use.
- 1 Not for people with galactosemia.

* NutraFlora® and scFOS® are not registered trademarks of Abbott Laboratories.

Ingredients

Liquid Unflavored:

Water, Corn Maltodextrin, Corn Syrup Solids, Sodium & Calcium Caseinates, Soy Protein Isolate, Canola Oil, Corn Oil, Fructooligosaccharides, Medium-Chain Triglycerides, Soy Fiber, Oat Fiber, Calcium Phosphate, Magnesium Phosphate, Potassium Citrate, Gum Arabic, Soy Lecithin, Potassium Chloride, Carboxymethylcellulose, Sodium Chloride, Ascorbic Acid, Choline Chloride, Sodium Citrate, L-Carnitine, Taurine, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Ferrous Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Beta-Carotene, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Cyanocobalamin, and Vitamin D3.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
53118	Jevity 1.2 Cal / 8-fl-oz (237-fl-oz) Can / 24 ct
62667	Jevity 1.2 Cal Ready-To-Hang / 1500-mL (1.6-QT) Bottle / 6 ct
62683	Jevity 1.2 Cal Ready-To-Hang / 1000-mL (1.1-QT) Bottle / 8 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)		1000 mL		1500 mL	
	Value	%RDI* / % DV	Value	%RDI* / % DV	Value	%RDI* / % DV
Nutrient Density, Cal/mL	1.2		1.2		1.2	
Protein, % Cal	18.5		18.5		18.5	
Carbohydrate, % Cal	52.5		52.5		52.5	
Fat, % Cal	29.0		29.0		29.0	
MCT:LCT	19:81		19:81		19:81	
Cal to Meet 100% RDIs	1200		1200		1200	
mL to Meet 100% RDIs	1000		1000		1000	
Total Cal:g Nitrogen	135:1		135:1		135:1	
Nonprotein Cal:g Nitrogen	110:1		110:1		110:1	
Osmolality, mOsm/kg H ₂ O	450		450		450	
Renal Solute Load, mOsm/L	465		465		465	
Viscosity	Nectar-Like (Room Temperature and Chilled)		Nectar-Like (Room Temperature and Chilled)		Nectar-Like (Room Temperature and Chilled)	
Minimum Tube Size for Gravity/Pump Feeding (Fr)	10/8		10/8		10/8	
Protein, g	13.2		55.5		83.3	
Fat, g	9.3		39.3		59.0	
Carbohydrate, g	40.2		169.4		254.0	
Dietary Fiber, g	4.3 [†]		18.0 [†]		27.0 [†]	
L-Carnitine, mg	36		150		225	
Taurine, mg	36		150		225	
Water, g/mL/cc	191		807		1211	
Calories	285		1200		1800	
Vitamin A, IU	1190 [‡]	24	5000 [‡]	100	7500 [‡]	150
Vitamin D, IU	96	24	400	100	600	150
Vitamin E, IU	11	37	45	150	68	225
Vitamin K, mcg	20	25	80	100	120	150
Vitamin C, mg	72	120	300	500	450	750
Folic Acid, mcg	145	36	600	150	900	225
Thiamin (Vitamin B1), mg	0.55	37	2.3	155	3.5	235
Riboflavin (Vitamin B2), mg	0.62	36	2.6	155	3.9	230
Vitamin B6, mg	0.72	36	3.0	150	4.5	225
Vitamin B12, mcg	2.2	37	9.0	150	14.0	235
Niacin, mg	7.2	36	30	150	45	225
Choline, mg	145	Not established	600	Not established	900	Not established
Biotin, mcg	110	37	450	150	675	225
Pantothenic Acid, mg	3.6	36	15	150	23	230

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)		1000 mL		1500 mL	
	Value	%RDI* / % DV	Value	%RDI* / % DV	Value	%RDI* / % DV
Sodium, mg	325	Not established	1350	Not established	2030	Not established
Sodium, mEq	14.1		58.7		88.3	
Potassium, mg	440	Not established	1850	Not established	2780	Not established
Potassium, mEq	11.3		47.4		71.3	
Chloride, mg	360	11	1500	44	2250	66
Chloride, mEq	10.2		42.4		63.6	
Calcium, mg	285	29	1200	120	1800	180
Phosphorus, mg	285	29	1200	120	1800	180
Magnesium, mg	96	24	400	100	600	150
Iodine, mcg	36	24	150	100	225	150
Manganese, mg	1.2	60	5.0	250	7.5	375
Copper, mg	0.48	24	2.0	100	3.0	150
Zinc, mg	5.5	37	23	155	34	225
Iron, mg	4.3	24	18	100	27	150
Selenium, mcg	17	24	70	100	105	150
Chromium, mcg	29	24	120	100	180	150
Molybdenum, mcg	36	48	150	200	225	300

Liquid Unflavored Footnotes & References

Per 8 fl oz (237 mL)

*RDI percentages for adults and children 4 or more years of age.

†

2.4 g of total dietary fiber from a unique fiber blend and 1.9 g of fructooligosaccharides.

‡795 IU of vitamin A activity supplied by 0.60 mg beta-carotene.

Per 1000 mL

*RDI percentages for adults and children 4 or more years of age.

†10.0 g of total dietary fiber from a unique fiber blend and 8.0 g of fructooligosaccharides.

‡3320 IU of vitamin A activity supplied by 2.5 mg beta-carotene.

Per 1500 mL

*RDI percentages for adults and children 4 or more years of age.

†15.0 g of total dietary fiber from a unique fiber blend and 12.0 g of fructooligosaccharides.

‡4980 IU of vitamin A activity supplied by 3.8 mg beta-carotene.

Preparation

Instructions for Use: Ready-To-Hang Container

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination. NOTE: Failure to follow the **Instructions for Use** increases the potential for microbial contamination and may reduce hangtime.

- 1 Pump feeding is recommended; use an 8 Fr or larger tube. If gravity feeding use a 10 Fr or larger tube.
- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Always follow directions for use provided with feeding set.

For Use with Enteral Feeding Set:

- 1 Remove the dust cover from the Abbott RTH Safety Screw Cap.
- 1 Remove the dust cover from the safety connector on the feeding set.
- 1 Insert the safety connector into the port on the Abbott RTH Safety Screw Cap, completely pierce foil.
- 1 Turn the safety connector clockwise until it is securely fastened.
- 1 Close clamp on set before inverting container.
- 1 Invert container and suspend, using the hanging ring on the bottom of the container.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.

1 **Instructions for Use: 8-fl-oz-can**

- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 Shake well. Jevity 1.2 is ready to use.
- 1 Clean top of container thoroughly before opening.
- 1 Once opened, cover, refrigerate and **use within 48 hours**.

Oral Feeding:

- 1 May be used for total or supplemental nutrition.
- 1 May be fed at room temperature or chilled.

Tube Feeding:

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Feed at room temperature.
- 1 Pump feeding is recommended; use an 8 Fr or larger tube. For gravity feeding use a 10 Fr or larger tube.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Avoid contamination during preparation and use.