

## Product Information: PediaSure® Peptide 1.0 Cal

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

# PediaSure® Peptide 1.0 Cal

## Peptide-Based Nutrition for Children

- 1 PediaSure Peptide 1.0 Cal is a nutritionally complete, peptide-based formula for the nutritional needs of children ages 1-13 years with malabsorption, maldigestion, and other GI conditions.
- 1 For oral or tube feeding.
- 1 For supplemental or sole-source nutrition.
- 1 Use under medical supervision.
- 1 Nutrient blend designed to help support tolerance and absorption:
  - Hydrolyzed whey-dominant protein.
  - Nutraflora® scFOS®\* prebiotic.
  - Structured lipids.
- 1 Nutrition to help support:
  - Absorption.
  - Tolerance.
  - Growth and development.
- 1 Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals for children 1 to 8 years of age in 1000 mL (1000 Cal); for children 9 to 13 years of age in 1500 mL (1500 Cal).
- 1 Only peptide-based pediatric product with Kosher status.
- 1 Halal.
- 1 Kosher.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.
- 1 Low-residue.



### Safety Precautions

- 1 Not intended for infants under 1 year of age unless specified by a physician.
- 1 Not for IV use.
- 1 Not for children with galactosemia.
- 1 **Feeding sets are for single-patient use only. Use clean technique to avoid set and/or product contamination.**

\* **NutraFlora® and scFOS® are not registered trademarks of Abbott Laboratories.**

### Ingredients

#### Liquid Unflavored:

Water, Corn Maltodextrin, Whey Protein Hydrolysate, Structured Lipid (Interesterified Canola, Medium-Chain Triglycerides), Hydrolyzed Sodium Caseinate, Medium-Chain Triglycerides, Canola Oil. Less than 0.5% of the Following: Short-Chain Fructooligosaccharides, Calcium Phosphate, Potassium Citrate, Cellulose Gel, Magnesium Phosphate, Soy Lecithin, Potassium Chloride, Magnesium Chloride, Ascorbic Acid, Choline Chloride, Cellulose Gum, m-Inositol, Carrageenan, Sodium Citrate, Taurine, Ferrous Sulfate, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, L-Carnitine, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Cupric Sulfate, Vitamin A Palmitate, Folic Acid, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate, Sodium Molybdate, Phylloquinone, Cyanocobalamin, and Vitamin D3.

**Allergens:** Contains milk and soy ingredients.

### Availability

List Number	Item
62119	PediaSure Peptide 1.0 Ready-to-Drink Vanilla / 8-fl-oz (237-mL) Bottle / 24 ct
62121	PediaSure Peptide 1.0 Ready-to-Drink Strawberry / 8-fl-oz (237-mL) Bottle / 24 ct
62123	PediaSure Peptide 1.0 Ready-to-Drink / 8-fl-oz (237-mL) Bottle / 24 ct
62729	PediaSure Peptide 1.0 Ready-To-Hang / 1000-mL (1.1-QT) Bottle / 8 ct

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)		1000 mL	
	Value	%RDI / %DV	Value	%RDI / %DV
Nutrient Density, Cal/mL	1.0		1.0	
Protein, % Cal	12		12	
Carbohydrate, % Cal	53		53	
Fat, % Cal	35		35	
MCT:LCT	50:50		50:50	
n6:n3	2.6:1		2.6:1	
mL to Meet Or Exceed 100% DRIs for Children 1-8 Years	1000		1000	
mL to Meet Or Exceed 100% DRIs for Children 9-13 Years	1500		1500	
Total Cal:g Nitrogen	208:1		208:1	
Nonprotein Cal:g Nitrogen	183:1		183:1	
Osmolality, mOsm/kg H2O	250		250	
Renal Solute Load, mOsm/L	293		293	
Viscosity	Thin (room temperature and chilled)		Thin (room temperature and chilled)	
Minimum Tube Size for Gravity/Pump Feeding (Fr)	10/8		10/8	
Protein, g	7.1		30.0	
Fat, g	9.6		40.5	
Trans Fat, g	0.0		0.0	
Carbohydrate, g	31.7		134	
Dietary Fiber, g	0.7		3.0	
scFOS, g	0.7		3.0	
L-Carnitine, mg	4.0		17	
Taurine, mg	18		76	
Water, g	200		845	
Calories	237		1000	
Vitamin A, IU	592		2500	
Vitamin D, IU	125		530	
Vitamin E, IU	5.4		23	
Vitamin K, mcg	14		59	
Vitamin C, mg	24		101	
Folic Acid, mcg	71		300	
Thiamin (Vitamin B1), mg	0.6		2.5	
Riboflavin (Vitamin B2), mg	0.5		2.1	
Vitamin B6, mg	0.6		2.5	
Vitamin B12, mcg	1.4		5.9	
Niacin, mg	2.4		10	
Choline, mg	71		300	
Biotin, mcg	45		190	
Pantothenic Acid, mg	2.4		10	

For more information, contact your Abbott Nutrition Representative or visit [www.abbottnutrition.com](http://www.abbottnutrition.com)

Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)		1000 mL	
	Value	%RDI / %DV	Value	%RDI / %DV
m-Inositol, mg	20		84	
Sodium, mg	170		717	
Sodium, mEq	7.4		31.2	
Potassium, mg	320		1350	
Potassium, mEq	8.2		34.6	
Chloride, mg	240		1010	
Chloride, mEq	6.8		28.5	
Calcium, mg	250		1060	
Phosphorus, mg	200		844	
Magnesium, mg	47		198	
Iodine, mcg	23		97	
Manganese, mg	0.4		1.7	
Copper, mg	0.2		1.0	
Zinc, mg	2.8		12	
Iron, mg	3.3		14	
Selenium, mcg	7.6		32	
Chromium, mcg	7.1		30	
Molybdenum, mcg	8.5		36	

For more information, contact your Abbott Nutrition Representative or visit [www.abbottnutrition.com](http://www.abbottnutrition.com)

### Preparation

#### Instructions for Use: 8-fl-oz Bottle

- 1 **Shake very well.**
- 1 Store unopened at room temperature.
- 1 After opening, refrigerate and **use within 48 hours.**

#### Tube Feeding:

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to child's condition and tolerance.
- 1 Additional fluid may be required.

#### Oral Feeding:

- 1 Best when chilled.

#### Instructions for Use: Ready-To-Hang Container

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination. NOTE: Failure to follow the **Instructions for Use** increases the potential for microbial contamination and reduces hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Always follow directions for use provided with feeding set.

#### For Use with Enteral Feeding Set:

- 1 Remove the dust cover from the Abbott RTH Safety Screw Cap.
- 1 Remove the dust cover from the safety connector on the feeding set.
- 1 Insert the safety connector into the port on the Abbott RTH Safety Screw Cap, completely pierce foil.
- 1 Turn the safety connector clockwise until it is securely fastened.
- 1 Close clamp on set before inverting container.
- 1 Invert container and suspend, using the hanging ring on the bottom of the container.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.