Premature Infant Formula with Iron

Product Information: Similac® Special Care® 20 With Iron

Premature Infant Formula with Iron

- 1 A 20 Cal/fl oz iron-fortified feeding for growing, low-birth-weight infants and premature infants.
- 1 Use under medical supervision.
- 1 A nutrient profile identical to Similac Special Care 24 per 100 Cal.
- 1 Has a unique blend of nucleotides to help support the developing immune system.
- 1 DHA and ARA, nutrients naturally found in breastmilk that are important for brain and eye development.
- 1 Lutein and DHA to support the developing eyes and brain.
- Calcium and phosphorus at concentrations and ratio (1.8:1) that supports intrauterine accretion rates.¹
- Approximately 2 mg iron/kg body weight per day, when fed at 120 Cal/kg body weight per day.
- 1 Mixes easily with other Similac® Special Care® formulas for a variety of high-caloric, nutrient-dense formulas—helps eliminate potential for mixing errors.
- 1 Kosher.
- 1 Halal.

Safety Precautions

- Not intended for feeding low-birth-weight infants after they reach a weight of 3600 g (approximately 8 lb) or as directed by a doctor.
- 1 If signs of intolerance develop, slow feeding or discontinue.
- 1 Never use a microwave oven to warm formula. Serious burns can result.

Ingredients

Liquid Unflavored:

Water, Nonfat Milk, Corn Syrup Solids, Medium-Chain Triglycerides, Lactose, Whey Protein Concentrate, Soy Oil, Coconut Oil. Less than 0.5% of the Following: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Calcium Phosphate, Ascorbic Acid, Magnesium Chloride, Soy Lecithin, Monoglycerides, Potassium Citrate, m-Inositol, Carrageenan, Calcium Carbonate, Sodium Citrate, Potassium Hydroxide, Ferrous Sulfate, Choline Bitartrate, Taurine, Niacinamide, Choline Chloride, L-Carnitine, Zinc Sulfate, Potassium Chloride, Salt, Potassium Phosphate, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Vitamin A Palmitate, Cupric Sulfate, Riboflavin, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Biotin, Phylloquinone, Sodium Selenate, Vitamin D3, Cyanocobalamin, and Nucleotides (Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate, Adenosine 5'-Monophosphate).

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
52418	Similac Special Care 20 with Iron Ready to Feed Inst / 2-fl-oz (59-mL) Bottle / 48 ct (discontinued)
56265	Similac Special Care 20 with Iron and Lutein & DHA RTF Inst / 2-fl-oz (59-mL) Bottle / 48 ct





¹ Mize et al. Am J Clin Nutr 1995;62:385-391.

Premature Infant Formula with Iron

Nutrition Information - Liquid Unflavored

Tradition information Elquid of	100 Cal	1000 mL
	100 Gai	1000 IIIL
	Value	Value
Calories	100	676
Volume, mL	148	1000
Protein, g	3.0	20.29
Fat, g	5.43	36.72
Carbohydrate, g	10.3	69.7
Water, g	133	899
Linoleic Acid, mg	700	4734
Vitamin A, IU	1250	8454
Vitamin D, IU	150	1014
Vitamin E, IU	4.0	27.1
Vitamin K, mcg	12	81.2
Thiamin (Vitamin B1), mcg	250	1691
Riboflavin (Vitamin B2), mcg	620	4193
Vitamin B6, mcg	250	1691
Vitamin B12, mcg	0.55	3.72
Niacin, mcg	5000	33815
Folic Acid (Folacin), mcg	37	250
Pantothenic Acid, mcg	1900	12850
Biotin, mcg	37	250
Vitamin C (Ascorbic Acid), mg	37	250
Choline, mg	10	68
Inositol, mg	40	271
Calcium, mg	180	1217
Phosphorus, mg	100	676
Magnesium, mg	12	81.2
Iron, mg	1.8	12.2
Zinc, mg	1.5	10.14
Manganese, mcg	12	81
Copper, mcg	250	1691
lodine, mcg	6	41
Selenium, mcg	1.8	12.2
Sodium, mg	43	291
Potassium, mg	129	872
Chloride, mg	81	548



Premature Infant Formula with Iron

Preparation

Ready to Feed

The health of the infant depends on carefully following the directions for preparation and use. Proper hygiene, handling and storage are important when preparing infant formula.

- Do not add water.
- 1 Shake very well.
- 1 Do not use if breakaway ring is missing or broken.
- 1 Twist off cap.
- 1 Twist on nipple and ring.
- 1 Pour desired amount into feeding container or attach nipple and ring (not included).
- 1 Once oral feeding begins, use within one hour or discard.
- 1 If signs of intolerance develop, slow feeding or discontinue.

Can be mixed to a variety of caloric densities using the Similac® Special Care® LiquiMix ® System. Follow physician's instructions.

Storage & Handling

Ready to Feed

- 1 Avoid prolonged exposure of bottles to light.
- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 If original bottle was not used for feeding, recap immediately and store in refrigerator for up to 24 hours.
- 1 Do not reuse bottle.

